

# F1 Issues

As of 5 January 2017  
Nuclear Regulation Authority (NRA), Japan

## Current Information on Radioactivity in Seawater

Sampling Date	Cs-134 (Bq/L)		Cs-137 (Bq/L)		H-3 (Bq/L)		*Gross Beta (Bq/L)	
	T-1	**T-2	T-1	**T-2	T-1	**T-2	T-1	**T-2
25 Dec. 2016	< 1.0	< 1.0	< 1.0	< 1.0	–	–	–	12
26 Dec. 2016	< 1.0	< 1.0	< 1.0	< 1.0	< 3.0	< 3.0	13	10
27 Dec. 2016	< 1.0	< 1.0	< 1.0	< 1.0	–	–	–	12
28 Dec. 2016	< 1.0	< 1.0	< 1.0	< 1.0	–	–	–	15
29 Dec. 2016	< 1.0	< 1.0	< 1.0	< 1.0	–	–	–	9.7
30 Dec. 2016	< 1.0	< 1.0	< 1.0	< 1.0	–	–	–	14
31 Dec. 2016	< 1.0	< 1.0	< 1.0	< 1.0	–	–	–	14
1 Jan. 2017	< 1.0	< 1.0	< 1.0	< 1.0	–	–	–	13
2 Jan. 2017	< 1.0	< 1.0	< 1.0	< 1.0	under analysis	under analysis	10	10


\*Gross Beta includes K-40 occurring naturally in seawater.

\*\*Sampling at T-2-1 was replaced by sampling at T-2 on/after 13 September 2016, because the road access to T-2-1 had been damaged by typhoon.

\*\*\*Under the Comprehensive Radiation Monitoring Plan, it is planned to conduct sampling at T-2-1. However, sampling at T-2-1 was replaced by sampling at T-2 on/after 13 September 2016, because the road access to T-2-1 had been damaged by typhoon.

Reference: "Comprehensive Radiation Monitoring Plan"

URL: <http://radioactivity.nsr.go.jp/en/list/274/list-1.html>



Concentrations of Cs-134, Cs-137, H-3 and Gross Beta remained low.

Details of past monitoring results (updated monthly) are open to the public via the following URL:

<http://radioactivity.nsr.go.jp/en/list/295/list-1.html>